arXiv:0901.1697v1 [math.NT] 13 Jan 2009

NOTE ON THE GENERALIZATION OF THE HIGHER ORDER q-GENOCCHI NUMBERS AND q-EULER NUMBERS

TAEKYUN KIM, YOUNG-HEE KIM, AND KYUNG-WON HWANG

Abstract Cangul-Ozden-Simsek[1] constructed the q-Genocchi numbers of high order using a fermionic p-adic integral on $\mathbb{Z}_p$, and gave Witt's formula and the interpolation functions of these numbers. In this paper, we present the generalization of the higher order q-Euler numbers and q-Genocchi numbers of Cangul-Ozden-Simsek. We define q-extensions of w-Euler numbers and polynomials, and w-Genocchi numbers and polynomials of high order using the multivariate fermionic p-adic integral on $\mathbb{Z}_p$. We have the interpolation functions of these numbers and polynomials. We obtain the distribution relations for q-extensions of w-Euler and w-Genocchi polynomials. We also have the interesting relation for q-extensions of these polynomials. We define (h, q)-extensions of w-Euler and w-Genocchi polynomials of high order. We have the interpolation functions for (h, q)-extensions of these polynomials. Moreover, we obtain some meaningful results of (h, q)-extensions of w-Euler and w-Genocchi polynomials.

2000 Mathematics Subject Classification : 11S80, 11B68

Key words and phrases : Genocchi numbers and polynomials, Euler numbers and polynomials, q-Genocchi numbers, q-Euler numbers, fermionic p-adic integral

1. Introduction, Definitions and Notations

Many authors have been studied on the multiple Genocchi and Euler numbers, and multiple zeta functions (cf. [1-2], [4-6], [9-10], [14], [17], [19], [22], [24]). In [10], Kim, the first author of this paper, presented a systematic study of some families of multiple q-Euler numbers and polynomials. By using the q-Volkenborn integration on $\mathbb{Z}_p$, Kim constructed the p-adic q-Euler numbers and polynomials of higher order, and gave the generating function of these numbers and the Euler q-ζ-function. In [14], Kim studied some families of multiple q-Genocchi and q-Euler numbers by using the multivariate p-adic q-Volkenborn integral on $\mathbb{Z}_p$, and gave interesting identities related to these numbers.

Recently, Cangul-Ozden-Simsek[1] constructed the q-Genocchi numbers of high order by using a fermionic p-adic integral on $\mathbb{Z}_p$, and gave Witt's formula and the interpolation functions of these numbers. In [17], Kim gave another constructions of the q-Euler and q-Genocchi numbers, which were different from those of Cangul-Ozden-Simsek. Kim obtained the interesting relationship between the q-w-Euler numbers and q-w-Genocchi numbers, and gave the interpolation functions of these numbers. In this paper, we will present the generalization of the higher order q-Euler numbers and q-Genocchi numbers of Cangul-Ozden-Simsek approaching as Kim did in [17].

Throughout this paper, let p be a fixed odd number and the symbols $\mathbb{Z}_p, \mathbb{Q}_p, \mathbb{C}$ and $\mathbb{C}_p$ denote the ring of p-adic rational integers, the field of p-adic rational numbers, the complex number field and the completion of algebraic closure of $\mathbb{Q}_p$, respectively. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = \frac{1}{p}$.

The symbol q can be treated as a complex number, $q \in \mathbb{C}$, or as a p-adic number, $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, then we always assume that $|q| < 1$. If $q \in \mathbb{C}_p$, then we usually assume that $|1-q|_p < 1$.

Now we will recall some q-notations. The q-basic natural numbers are defined by $[n]_q = \frac{1-q^n}{1-q} = 1 + q + q^2 + \cdots + q^{n-1}$ $(n \in \mathbb{N})$, $[n]_{-q} = \frac{1-(-q)^n}{1+q}$ and the q-factorial by $[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$. In this paper, we use the notation $[x]_q = \frac{1-q^x}{1-q}$ and $[x]_{-q} = \frac{1-(-q)^x}{1+q}$. Hence $\lim_{q\to 1}[x]_q = x$ for any x with $|x|_p \le 1$ in the present p-adic case (see [1-25]).

The q-shift factorial is given by

$$(a:q)_0 = 1, \quad (a:q)_k = (1-a)(1-aq)\cdots(1-aq^{k-1}).$$

We note that $\lim_{q\to 1}(a:q)_k = (1-a)^k$. It is known that

$$(a:q)_\infty = (1-a)(1-aq)(1-aq^2)\cdots = \prod_{i=1}^{\infty}(1-aq^{i-1}), \quad \text{(see [8])}.$$

From the definition of the q-shift factorial, we note that

$$(a:q)_k = \frac{(a:q)_\infty}{(aq^k:q)_\infty}.$$

Since $\binom{-\alpha}{l} = (-1)^l\binom{\alpha+l-1}{l}$, it follows that

$$\frac{1}{(1-z)^\alpha} = (1-z)^{-\alpha} = \sum_{l=0}^{\infty}\binom{-\alpha}{l}(-z)^l = \sum_{l=0}^{\infty}\binom{\alpha+l-1}{l}z^l.$$

The q-binomial theorem is given by

$$\sum_{n=0}^{\infty}\frac{(a:q)_n}{(q:q)_n}z^n = \frac{(az:q)_\infty}{(z:q)_\infty},$$

where $z, q \in \mathbb{C}$ with $|z| < 1$, $|q| < 1$. For the special case, when $a = q^\alpha (\alpha \in \mathbb{C})$, we can write as follows:

$$\begin{aligned}
\frac{1}{(z:q)_\alpha} &= \frac{(zq^\alpha:q)_\infty}{(z:q)_\infty} = \sum_{n=0}^{\infty}\frac{(q^\alpha:q)_n}{(q:q)_n}z^n \\
&= \sum_{n=0}^{\infty}\frac{(1-q^\alpha)(1-q^{\alpha+1})\cdots(1-q^{\alpha+n-1})}{(1-q)(1-q^2)\cdots(1-q^n)}z^n \\
&= \sum_{n=0}^{\infty}\frac{[\alpha]_q[\alpha+1]_q\cdots[\alpha+n-1]_q}{[1]_q[2]_q\cdots[n]_q}z^n \\
&= \sum_{n=0}^{\infty}\frac{[\alpha]_q[\alpha+1]_q\cdots[\alpha+n-1]_q}{[n]_q!}z^n.
\end{aligned}$$

The q-binomial coefficients are defined by

$$\binom{n}{k}_q = \frac{[n]_q!}{[k]_q![n-k]_q!} = \frac{[n]_q[n-1]_q\cdots[n-k+1]_q}{[k]_q!}, \quad \text{(see [14], [16])}.$$

Hence it follows that

$$\frac{1}{(z:q)_\alpha}=\sum_{n=0}^{\infty}\binom{n+\alpha-1}{n}_q z^n,$$

which converges to $\frac{1}{(1-z)^\alpha}=\sum\limits_{n=0}^{\infty}\binom{n+\alpha-1}{n}z^n$ as $q\to 1$.

We say that f is a uniformly differentiable function at a point $a\in\mathbb{Z}_p$, and write $f\in UD(\mathbb{Z}_p)$, the set of uniformly differentiable function, if the difference quotients $F_g(x,y)=\frac{f(x)-f(y)}{x-y}$ have a limit $l=f'(a)$ as $(x,y)\to(a,a)$. For $f\in UD(\mathbb{Z}_p)$, the q-deformed bosonic p-adic integral is defined as

$$I_q(f)=\int_{\mathbb{Z}_p}f(x)d\mu_q(x)=\lim_{N\to\infty}\sum_{x=0}^{p^N-1}f(x)\frac{q^x}{[p^N]_q},$$

and the q-deformed fermonic p-adic integral is defined by

$$I_{-q}(f)=\int_{\mathbb{Z}_p}f(x)d\mu_{-q}(x)=\lim_{N\to\infty}\sum_{x=0}^{p^N-1}f(x)\frac{(-q)^x}{[p^N]_{-q}}.$$

The fermionic p-adic integral on $\mathbb{Z}_p$ is defined as

$$I_{-1}(f)=\lim_{q\to1}I_{-q}(f)=\int_{\mathbb{Z}_p}f(x)d\mu_{-1}(x).$$

It follows that $I_{-1}(f_1)=-I_{-1}(f)+2f(0)$, where $f_1(x)=f(x+1)$. For details, see [4-17].

The classical Euler polynomials $E_n(x)$ are defined as

$$\frac{2}{e^t+1}e^{xt}=\sum_{x=0}^{\infty}E_n(x)\frac{t^n}{n!},$$

and the Euler numbers E_n are defined as $E_n=E_n(0)$, (see [1-25]). The Genocchi numbers are defined as

$$\frac{2t}{e^t+1}=\sum_{n=0}^{\infty}G_n\frac{t^n}{n!}\quad\text{for}\quad |t|<\pi,$$

and the Genocchi polynomials $G_n(x)$ are defined as

$$\frac{2t}{e^t+1}e^{xt}=\sum_{n=0}^{\infty}G_n(x)\frac{t^n}{n!},\quad\text{(see [12], [14], [21]).}$$

It is known that the w-Euler polynomials $E_{n,w}(x)$ are defined as

$$\frac{2}{we^t+1}e^{xt}=\sum_{x=0}^{\infty}E_{n,w}(x)\frac{t^n}{n!},$$

and $E_{n,w}=E_{n,w}(0)$ are called the w-Euler numbers. The w-Genocchi polynomials $G_{n,w}(x)$ are defined as

$$\frac{2t}{we^t+1}e^{xt}=\sum_{x=0}^{\infty}G_{n,w}(x)\frac{t^n}{n!},$$

and $G_{n,w}=G_{n,w}(0)$ are called the w-Genocchi numbers, (see [1]).

The w-Euler polynomials $E_{n,w}^{(r)}(x)$ of order r are defined as

$$(\frac{2}{we^t+1})^r e^{xt} = \sum_{x=0}^{\infty} E_{n,w}^{(r)}(x)\frac{t^n}{n!}, \quad \text{(see [1])},$$

and $E_{n,w}^{(r)} = E_{n,w}^{(r)}(0)$ are called the w-Euler numbers of order r. The w-Genocchi polynomials $G_{n,w}^{(r)}(x)$ of order r are defined as

$$\frac{2t}{we^t+1}e^{xt} = \sum_{x=0}^{\infty} G_{n,w}^{(r)}(x)\frac{t^n}{n!}, \quad \text{(see [1])},$$

and $G_{n,w}^{(r)} = G_{n,w}^{(r)}(0)$ are called the w-Euler numbers of order r. When $r = 1$ and $w = 1$, $E_{n,w}^{(r)}(x)$ and $E_{n,w}^{(r)}$ are the ordinary Euler polynomials and numbers, and $G_{n,w}^{(r)}(x)$ and $G_{n,w}^{(r)}$ are the ordinary Genocchi polynomials and numbers, respectively.

In Section 2, we define q-extensions of w-Euler numbers and polynomials of order r and w-Genocchi numbers and polynomials of order r, respectively, using the multivariate fermionic p-adic integral on $\mathbb{Z}_p$. We obtain the interpolation functions of these numbers and polynomials. We have the distribution relations for q-extensions of w-Euler polynomials and those of w-Genocchi polynomials. We obtain the interesting relation for q-extensions of these polynomials. We also define (h,q)-extensions of w-Euler and w-Genocchi polynomials of order r. We have the interpolation functions for (h,q)-extensions of these polynomials. Moreover, we obtain some meaningful results of (h,q)-extensions of w-Euler and w-Genocchi polynomials when $h = r-1$.

2. On the extension of the higher order q-Genocchi numbers and q-Euler numbers of Cangul-Ozden-Simsek

In this section, we assume that $w \in \mathbb{C}_p$ with $|1-w|_p < 1$ and $q \in \mathbb{C}_p$ with $|1-q|_p < 1$. Recently, Cangul-Ozden-Simsek[1] constructed w-Genocchi numbers of order r, $G_{n,w}^{(r)}$, as follows.

$$\begin{aligned} t^r \int_{\mathbb{Z}_p^r} & w^{x_1+\cdots+x_r} e^{t(x_1+\cdots+x_r)} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \\ &= 2^r(\frac{t}{we^t+1})^r = \sum_{n=0}^{\infty} G_{n,w}^{(r)}\frac{t^n}{n!}, \end{aligned} \tag{1}$$

where $\int_{\mathbb{Z}_p^r} = \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}$ ($r-$times) and $r \in \mathbb{N}$. They also consider the q-extension of $G_{n,w}^{(r)}$ as follows.

$$\begin{aligned} t^r \int_{\mathbb{Z}_p^r} & q^{\sum_{i=1}^{r}(h-i+1)x_i} e^{t(\sum_{i=1}^{r} x_i)} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \\ &= \frac{2^r t^r}{(q^h e^t+1)\cdots(q^{h-r+1}e^t+1)} = \sum_{n=0}^{\infty} G_{n,q}^{(h,r)}\frac{t^w}{n!}. \end{aligned} \tag{2}$$

From (2), they obtained the following interesting formula:

$$G_{n+r,q}^{(r-1,r)} = 2^r r! \binom{n+r}{r} \sum_{v=0}^{\infty} \binom{r+v-1}{v}_q (-1)^v v^n. \tag{3}$$

It seems to be interested to find the numbers corresponding to

$$2^r r! \binom{n+r}{r} \sum_{v=0}^{\infty} \binom{r+v-1}{v}_q (-1)^v [v]_q^n.$$

In the viewpoint of the q-extension of (1) using the multivariate $p-$adic integral on $\mathbb{Z}_p$, we define the q-analogue of w-Euler numbers of order r, $E_{n,w,q}^{(r)}$, as follows.

$$E_{n,w,q}^{(r)} = \int_{\mathbb{Z}_p^r} w^{x_1+\cdots+x_r} [x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r). \tag{4}$$

From (4), we note that

$$\begin{aligned} E_{n,w,q}^{(r)} &= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} (-1)^l (\frac{1}{1+qw})^r \\ &= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} (-1)^l \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m q^{lm} w^m \\ &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m w^m [m]_q^n. \end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 1. *Let $r \in \mathbb{N}$ and $n \in \mathbb{Z}_+$. Then we have*

$$E_{n,w,q}^{(r)} = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m w^m [m]_q^n. \tag{5}$$

Let $F^{(r)}(t,w|q) = \sum_{n=0}^{\infty} E_{n,w,q}^{(r)} \frac{t^n}{n!}$. By (4) and (5), we see that

$$\begin{aligned} F^{(r)}(t,w|q) &= \int_{\mathbb{Z}_p^r} w^{x_1+\cdots+x_r} e^{t[x_1+\cdots+x_r]_q} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \\ &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m w^m e^{t[m]_q}. \end{aligned}$$

Thus we obtain the following corollary.

Corollary 2. *Let $F^{(r)}(t,w|q) = \sum_{n=0}^{\infty} E_{n,w,q}^{(r)} \frac{t^n}{n!}$. Then we have*

$$F^{(r)}(t,w|q) = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m w^m e^{t[m]_q}.$$

Let us define the q-extension of w-Euler polynomials of order r as follows.

$$E_{n,w,q}^{(r)}(x) = \int_{\mathbb{Z}_p^r} w^{x_1+\cdots+x_r} [x+x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r). \tag{6}$$

By (6), we have that

$$\begin{aligned} E_{n,w,q}^{(r)}(x) &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l q^{lx}(\frac{1}{1+q^l w})^r \\ &= 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}(-1)^m w^m [m+x]_q^n \end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 3. *Let $r \in \mathbb{N}$ and $n \in \mathbb{Z}_+$. Then we have*

$$E_{n,w,q}^{(r)}(x) = 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}(-1)^m w^m [m+x]_q^n. \tag{7}$$

Let $F^{(r)}(t,w,x|q) = \sum_{n=0}^{\infty} E_{n,w,q}^{(r)}(x)\frac{t^n}{n!}$. By (6) and (7), we have

$$\begin{aligned} F^{(r)}(t,w,x|q) &= \int_{\mathbb{Z}_p^r} w^{x_1+\cdots+x_r} e^{t[x+x_1+\cdots+x_r]_q} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \\ &= 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}(-1)^m w^m e^{t[m+x]_q}. \end{aligned}$$

Therefore we have the following corollary.

Corollary 4. *Let $F^{(r)}(t,w,x|q) = \sum_{n=0}^{\infty} E_{n,w,q}^{(r)}(x)\frac{t^n}{n!}$. Then we have*

$$F^{(r)}(t,w,x|q) = 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}(-1)^m w^m e^{t[m+x]_q}. \tag{8}$$

Now we define the q-extension of w-Genocchi polynomials of order r, $G_{n,w,q}^{(r)}(x)$, as follows.

$$2^r t^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}(-1)^m w^m e^{t[m+x]_q} = \sum_{n=0}^{\infty} G_{n,w,q}^{(r)}(x)\frac{t^n}{n!}. \tag{9}$$

Then we have

$$\begin{aligned} (10)\quad & \sum_{n=0}^{\infty} G_{n,w,q}^{(r)}(x)\frac{t^n}{n!} \\ = \quad & t^r\int_{\mathbb{Z}_p^r} w^{x_1+\cdots+x_r} e^{t[x+x_1+\cdots+x_r]_q} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \\ = \quad & \sum_{n=0}^{\infty}\int_{\mathbb{Z}_p^r} w^{x_1+\cdots+x_r}[x+x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) r!\binom{n+r}{r}\frac{t^{n+r}}{(n+r)!}. \end{aligned}$$

By comparing the coefficients on the both sides of (10), we see that

$$G_{0,w,q}^{(r)}(x) = G_{1,w,q}^{(r)}(x) = \cdots = G_{r-1,w,q}^{(r)}(x) = 0,$$

and

$$\begin{aligned}(11)\qquad & G^{(r)}_{n+r,w,q}(x)\\ = \ & r!\binom{n+r}{r}\int_{\mathbb{Z}_p^r} w^{x_1+x_2+\cdots+x_r}[x+x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)\\ = \ & r!\binom{n+r}{r}E^{(r)}_{n,w,q}(x).\end{aligned}$$

In the special case of $x=0$, $G^{(r)}_{n,w,q}(0)=G^{(r)}_{n,w,q}$ are called the q-extension of w-Genocchi numbers of order r. By (11), we have the following theorem.

Theorem 5. *Let* $r\in\mathbb{N}$ *and* $n\in\mathbb{Z}_+$. *Then we have*

$$\begin{aligned}\frac{G^{(r)}_{n+r,w,q}(x)}{r!\binom{n+r}{r}} &= \int_{\mathbb{Z}_p^r} w^{x_1+\cdots+x_r}[x+x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)\\ &= E^{(r)}_{n,w,q}(x),\end{aligned}$$

and $G^{(r)}_{0,w,q}(x)=G^{(r)}_{1,w,q}(x)=\cdots=G^{(r)}_{r-1,w,q}(x)=0.$

Now we consider the distribution relation for the q-extension of w-Euler polynomials of order r. For $d\in\mathbb{N}$ with $d\equiv 1 \pmod 2$, by (8), we see that

$$\begin{aligned}(12)\quad & F^{(r)}(t,w,x|q)\\ = \ & 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}(-1)^m w^m e^{t[m+x]_q}\\ = \ & \sum_{a_1,\cdots a_r=0}^{d-1}(\prod_{i=1}^{r} w^{a_i})(-1)^{a_1+\cdots+a_r}2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}(-1)^m w^{md}e^{t[d]_q[m+\frac{a_1+\cdots+a_r+x}{d}]_{q^d}}\\ = \ & \sum_{a_1,\cdots a_r=0}^{d-1}(\prod_{i=1}^{r} w^{a_i})(-1)^{a_1+\cdots+a_r}F^{(r)}([d]_q t,\, w^d,\, \frac{a_1+\cdots+a_r+x}{d}\,|\, q^d).\end{aligned}$$

By (12), we obtain the following distribution relations for $E^{(r)}_{n,w,q}(x)$ and $G^{(r)}_{n+r,w,q}(x)$, respectively.

Theorem 6. *Let* $r\in\mathbb{N}$, $n\in\mathbb{Z}_+$ *and* $d\in\mathbb{N}$ *with* $d\equiv 1$ (*mod* 2). *Then we have*

$$E^{(r)}_{n,w,q}(x)=[d]_q^n\sum_{a_1,\cdots a_r=0}^{d-1}(\prod_{i=1}^{r} w^{a_i})(-1)^{a_1+\cdots+a_r}E^{(r)}_{n,w^d,q^d}(\frac{a_1+\cdots+a_r+x}{d}).$$

Furthermore,

$$G^{(r)}_{n+r,w,q}(x)=[d]_q^n\sum_{a_1,\cdots a_r=0}^{d-1}(\prod_{i=1}^{r} w^{a_i})(-1)^{a_1+\cdots+a_r}G^{(r)}_{n+r,w^d,q^d}(\frac{a_1+\cdots+a_r+x}{d}).$$

For the extension of (2), we consider the (h,q)-extension of w-Euler polynomials of order r. For $h\in\mathbb{Z}$, $r\in\mathbb{N}$ and $n\in\mathbb{Z}_+$, let us define the (h,q)-extension of w-Euler polynomial of order r as follows.

$$\begin{aligned} & E^{(h,r)}_{n,w,q}(x) \\ &= \int_{\mathbb{Z}_p^r} w^{x_1+\cdots+x_r}[x+x_1+\cdots+x_r]_q^n q^{\sum_{i=1}^{r}(h-i+1)x_i} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r). \end{aligned} \tag{13}$$

From (13), we note that

$$\begin{aligned} E^{(h,r)}_{n,w,q}(x) &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\frac{\binom{n}{l}(-1)^l q^{lx}}{(1+q^{l+h}w)(1+q^{l+h-1}w)\cdots(1+q^{l+h-r+1}w)} \\ &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\frac{\binom{n}{l}(-1)^l q^{lx}}{(-q^{l+h}w : q^{-1})_r} \qquad (14) \\ &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l q^{lx}\sum_{m=0}^{\infty}\binom{m+r-1}{m}_{q^{-1}}(-1)^m q^{(l+h)m}w^m \\ &= 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}_{q^{-1}}(-1)^m q^{hm}w^m[m+x]_q^n. \end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 7. *Let $h \in \mathbb{Z}$, $r \in \mathbb{N}$ and $n \in \mathbb{Z}_+$. Then we have*

$$\begin{aligned} E^{(h,r)}_{n,w,q}(x) &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\frac{\binom{n}{l}(-1)^l q^{lx}}{(-q^{l+h}w : q^{-1})_r} \\ &= 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}_{q^{-1}}(-1)^m q^{hm}w^m[m+x]_q^n. \end{aligned} \tag{15}$$

We also have the following result.

Corollary 8. *Let $F^{(h,r)}(t,w,x|q) = \sum_{n=0}^{\infty} E^{(h,r)}_{n,w,q}(x)\frac{t^n}{n!}$. Then we have*

$$F^{(h,r)}(t,w,x|q) = 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}_{q^{-1}}(-1)^m q^{hm}w^m e^{t[m+x]_q}. \tag{16}$$

Remark 1. *In the special case $x = 0$, $E^{(h,r)}_{n,w,q}(0) = E^{(h,r)}_{n,w,q}$ are called the (h,q)-extension of w-Euler numbers of order r.*

If we take $h = r-1$ in (14), then we have

$$\begin{aligned} E^{(r-1,r)}_{n,w,q}(x) &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\frac{\binom{n}{l}(-1)^l q^{lx}}{(1+q^{l+r-1}w)(1+q^{l+r-2}w)\cdots(1+q^{l}w)} \\ &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\frac{\binom{n}{l}(-1)^l q^{lx}}{(-q^{l}w : q)_r} \qquad (17) \\ &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l q^{lx}\sum_{m=0}^{\infty}\binom{m+r-1}{m}_{q}(-1)^m q^{lm}w^m \\ &= 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}_{q}(-1)^m w^m[m+x]_q^n. \end{aligned}$$

Then we have the following theorem.

Theorem 9. *Let* $r \in \mathbb{N}$ *and* $n \in \mathbb{Z}_+$. *Then we have*

$$\begin{aligned} E_{n,w,q}^{(r-1,r)}(x) &= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \frac{\binom{n}{l}(-1)^l q^{lx}}{(-q^l w : q)_r} \\ &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m w^m [m+x]_q^n. \end{aligned}$$

We also the following corollary.

Corollary 10. *Let* $F^{(r-1,r)}(t,w,x|q) = \sum_{n=0}^{\infty} E_{n,w,q}^{(r-1,r)}(x) \frac{t^n}{n!}$. *Then we have*

$$F^{(r-1,r)}(t,w,x|q) = 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m w^m e^{t[m+x]_q}. \tag{18}$$

From (18), we note that

$$\begin{aligned} F^{(r-1,r)}(t,w,x|q) &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m w^m e^{t[m+x]_q} \\ &= \sum_{a_1,\cdots a_r=0}^{d-1} q^{\sum_{i=0}^{r}(r-i)a_i} (-1)^{a_1+\cdots+a_r} w^{a_1+\cdots+a_r} \\ &\quad \times 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_{q^d} (-1)^m w^{md} e^{t[d]_q[m+\frac{a_1+\cdots+a_r+x}{d}]_{q^d}} \qquad (19) \\ &= \sum_{a_1,\cdots a_r=0}^{d-1} q^{\sum_{i=0}^{r}(r-i)a_i} (-1)^{a_1+\cdots+a_r} w^{a_1+\cdots+a_r} \\ &\quad \times F^{(r-1,r)}([d]_q t, w^d, \frac{a_1+\cdots+a_r+x}{d} | q^d), \end{aligned}$$

where $d \in \mathbb{N}$ with $d \equiv 1 \pmod 2$. By (19), we obtain the following the distribution relation for $E_{n,w,q}^{(r-1,r)}(x)$.

Theorem 11. *For* $r \in \mathbb{N}$, $n \in \mathbb{Z}_+$ *and* $d \in \mathbb{N}$ *with* $d \equiv 1$ *(mod 2). Then we have*

$$\begin{aligned} & E_{n,w,q}^{(r-1,r)}(x) \\ &= [d]_q^n \sum_{a_1,\cdots a_r=0}^{d-1} q^{\sum_{i=0}^{\infty}(r-i)a_i} (-1)^{a_1+\cdots+a_r} w^{a_1+\cdots+a_r} E_{n,w^d,q^d}^{(r-1,r)}(\frac{a_1+\cdots+a_r+x}{d}). \end{aligned}$$

Now we define the (h,q)-extension of w-Genocchi polynomials $G_{n,w,q}^{(h,r)}(x)$ of order r as follows.

$$2^r t^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_{q^{-1}} (-1)^m q^{hm} w^m e^{t[m+x]_q} = \sum_{n=0}^{\infty} G_{n,w,q}^{(h,r)}(x) \frac{t^n}{n!}. \tag{20}$$

Then we have

$$
\begin{aligned}
&\sum_{n=0}^{\infty} G_{n,w,q}^{(h,r)}(x)\frac{t^n}{n!} \\
&= t^r \int_{\mathbb{Z}_p^r} q^{\sum_{i=0}^{\infty}(h-i+1)x_i} w^{x_1+\cdots+x_r} e^{t[x+x_1+\cdots+x_r]_q} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \qquad (21)\\
&= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p^r} q^{\sum_{i=0}^{\infty}(h-i+1)x_i} w^{x_1+\cdots+x_r} [x+x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \\
&\quad \times r!\binom{n+r}{r}\frac{t^{n+r}}{(n+r)!}.
\end{aligned}
$$

From (13) and (21), we derive the following result.

Theorem 12. *Let $r \in \mathbb{N}$ and $n \in \mathbb{Z}_+$. Then we have*

$$
\begin{aligned}
\frac{G_{n+r,w,q}^{(h,r)}(x)}{r!\binom{n+r}{r}} &= \int_{\mathbb{Z}_p^r} q^{\sum_{i=0}^{\infty}(h-i+1)x_i} w^{x_1+\cdots+x_r} [x+x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \\
&= E_{n,w,q}^{(h,r)}(x),
\end{aligned}
$$

and $G_{0,w,q}^{(h,r)}(x) = G_{1,w,q}^{(h,r)}(x) = \cdots = G_{r-1,w,q}^{(h,r)}(x) = 0.$

When $h = r-1$ in Theorem 12, we have

$$
\begin{aligned}
\frac{G_{n+r,w,q}^{(r-1,r)}(x)}{r!\binom{n+r}{r}} &= \int_{\mathbb{Z}_p^r} q^{\sum_{i=0}^{\infty}(r-i)x_i} w^{x_1+\cdots+x_r} [x+x_1+\cdots+x_r]_q^n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \\
&= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m w^m [m+x]_q^n \\
&= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n} \frac{\binom{n}{l}(-1)^l q^{lx}}{(-q^l w : q)_r} \\
&= E_{n,w,q}^{(r-1,r)}(x).
\end{aligned}
$$

Remark 2. *In the special case $x = 0$, $G_{n,w,q}^{(h,r)}(0) = G_{n,w,q}^{(h,r)}$ are called the (h,q)-extension of w-Genocchi numbers of order r.*

References

[1] I. N. Cangul, H. Ozden, Y. Simsek, *A new approach to q-Genocchi numbers and their interpolations*, Nonlinear Analysis (2008), doi:10.1016/j.na.2008.11.040.

[2] M. Cenkci, Y. Simsek, V. Kurt, *Multiple two-variable p-adic q-L-function and its behavior at $s = 0$*, Russ. J. Math. Phys. **15** (2008) no. 4, 447-459.

[3] G. Kim, B. Kim, J. Choi, *The DC algorithm for computing sums of powers of consecutive integers and Bernoulli numbers*, Adv. Stud. Contemp. Math. (Kyungshang) **17** (2008), no. 2, 137–145.

[4] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288–299.

[5] T. Kim, *On Euler-Barnes multiple zeta functions*, Russ. J. Math. Phys. **10** (2003), no. 3, 261–267.

[6] T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russ. J. Math. Phys. **11** (2004), no. 12, 71–76.

- [7] T. Kim, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russ. J. Math. Phys. **12** (2005), no. 2, 186–196.
- [8] T. Kim, *q-Generalized Euler numbers and polynomials*, Russ. J. Math. Phys. **13** (2006), no. 3, 293–298.
- [9] T. Kim, *Multiple p-adic L-function*, Russ. J. Math. Phys. **13** (2006), no. 2, 151–157.
- [10] T. Kim, *q-Euler numbers and polynomials associated with p-adic q-integrals*, J. Nonlinear Math. Phys. **14** (2007), no. 1, 15–27.
- [11] T. Kim, *A note on p-adic q-integral on $\mathbb{Z}_p$ associated with q-Euler numbers*, Adv. Stud. Contemp. Math. (Kyungshang) **15** (2007), 133–138.
- [12] T. Kim, *On the q-extension of Euler and Genocchi numbers*, J. Math. Anal. Appl. **326** (2007), 1458–1465.
- [13] T. Kim, *q-Extension of the Euler formula and trigonometric functions*, Russ. J. Math. Phys. **14** (2007), no. 3, 275–278.
- [14] T. Kim, *On the multiple q-Genocchi and Euler numbers*, Russ. J. Math. Phys. **15** (2008) no.4, 481-486.
- [15] T. Kim, *The modified q-Euler numbers and polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **16** (2008), 161–170.
- [16] T. Kim, *q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russ. J. Math. Phys. **15** (2008), no.1, 51-57.
- [17] T. Kim, *New approach to q-Euler, Genocchi numbers and their interpolation functions*, arXiv:0901.0353v1 [math.NT].
- [18] Y.-H. Kim, W. Kim, L.-C. Jang, *On the q-extension of Apostol-Euler numbers and polynomials*, Abstr. Appl. Anal. **2008** (2008), Article ID 296159, 10 pages.
- [19] H. Ozden, I. N. Cangul, Y. Simsek, *Multivariate interpolation functions of higher-order q-Euler numbers and their applications*, Abstr. Appl. Anal. **2008** (2008), Art. ID 390857, 16 pages.
- [20] H. Ozden, Y. Simsek, *A new extension of q-Euler numbers and polynomials related to their interpolation functions*, Appl. Math. Lett. **21** (2008), 934–939.
- [21] K. H. Park, Y.-H. Kim, *On some arithmetical properties of the Genocchi numbers and polynomials*, Advances in Difference Equations (2008), http://www.hindawi.com/journals/ade/aip.195049.html.
- [22] Y. Simsek, *Complete sum of products of (h, q)-extension of the Euler polynomials and numbers*, arXiv:0707.2849v1 [math.NT].
- [23] J. V. Leyendekkers, A. G. Shannon, C. K. Wong, *Integer structure analysis of the product of adjacent integers and Euler's extension of Fermat's last theorem*, Adv. Stud. Contemp. Math. (Kyungshang) **17** (2008), no. 2, 221–229.
- [24] H.M. Srivastava, T. Kim, Y. Simsek, *q-Bernoulli numbers and polynomials associated with multiple q-zeta functions and basic L-series*, Russ. J. Math. Phys. **12** (2005), no. 2, 241–268.
- [25] Z. Zhang, Y. Zhang, *Summation formulas of q-series by modified Abel's lemma*, Adv. Stud. Contemp. Math. (Kyungshang) **17** (2008), no. 2, 119–129.

Taekyun Kim. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `tkkim@kw.ac.kr`

Young-Hee Kim. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `yhkim@kw.ac.kr`

Kyung-Won Hwang. Department of General education, Kookmin university, Seoul 136-702, Republic of Korea,
E-mail address: `khwang7@kookmin.ac.kr`